

# HEXYL AMINOLEVULINATE FLUORESCENCE CYSTOSCOPY: A NEW DIAGNOSTIC TOOL FOR THE PHOTODIAGNOSIS OF SUPERFICIAL BLADDER CANCER – A MULTICENTER STUDY



## **Authors:**

Patrice Jichlinski, Louis Guillou, Steinar Karlsen et al

## **Citation:**

Journal of Urology 2003; 170: 226-9.



To examine the sensitivity and specificity of Hexvix cystoscopy in patients with superficial bladder cancer

## METHODS


- In this open study 52 patients with suspected superficial bladder cancer were investigated at 4 European centers
- 50ml Hexvix® (8mM) was instilled into the bladder of patients 1 hour prior to cystoscopy and transurethral resection of the bladder
- The bladder was first inspected under white light and the number and location of tumors noted. The light system was then changed to fluorescence mode (blue light) and all fluorescing lesions documented
- At the end of the fluorescence inspection, 5 random biopsies per patient were taken from areas of the urothelium that appeared normal under white light
- Following this, all lesions mapped in white light were biopsied or resected, followed by the same procedure with additional fluorescing lesions. Tissue from all biopsies and resections was collected for histology



## RESULTS

- Hexvix® blue-light cystoscopy was highly efficient at detecting tumors compared with standard white light cystoscopy

### PATIENT DETECTION RATE



- Hexvix® blue-light cystoscopy detected 30% more bladder cancer patients than standard white light cystoscopy
- The 10 cases missed by white light inspection but detected with Hexvix® blue-light cystoscopy were CIS tumors
- No local or systemic adverse events clearly related to the use of Hexvix® blue-light cystoscopy were reported

## CONCLUSION

Hexvix® blue-light cystoscopy is a sensitive, well-tolerated diagnostic procedure for bladder cancer that showed improved detection of bladder tumors, in particular CIS

The procedure can easily be implemented in current clinical practice

# PRESCRIBING INFORMATION HEXVIX (HEXAMINOLEVULINATE)

Please refer to full national Summary of Product Characteristics (SPC) before prescribing. Indications and approvals may vary in different countries. Further information available on request. Hexvix 85 mg, powder and solvent for solution for intravesical use.

**PRESENTATION** Pack of one 10ml glass vial containing 85mg of hexaminolevulinate as 100mg hexaminolevulinate hydrochloride as a powder and one 50ml polypropylene vial containing solvent. After reconstitution in 50ml of solvent, 1ml of the solution contains 1.7mg hexaminolevulinate which corresponds to a 8mmol/l solution of hexaminolevulinate.

**INDICATIONS** This medicinal product is for diagnostic use only. Hexvix blue light fluorescence cystoscopy is indicated as adjunct to standard white light cystoscopy to contribute to the diagnosis and management of bladder cancer in patients with known or high suspicion of bladder cancer.

**DOSAGE AND METHOD OF ADMINISTRATION** Hexvix cystoscopy should only be performed by health care professionals trained specifically in Hexvix cystoscopy. The bladder should be drained before the instillation. Adults (including the elderly): 50 ml of 8 mmol/l reconstituted solution (see section 6.6) is instilled into the bladder through a catheter. The patient should retain the fluid for approximately 60 minutes. Following evacuation of the bladder, the cystoscopic examination in blue light should start within approximately 60 minutes. The cystoscopic examination should not be performed more than 3 hours after Hexvix is instilled in the bladder. Also if the retention time in the bladder is considerable shorter than one hour, examination should start no earlier than after 60 minutes. No minimum retention time has been identified making examination non-informative. For optimal visualisation it is recommended to examine and map the entire bladder under both white and blue light before any surgical measures are initiated. Biopsies of all mapped lesions should normally be taken under white light and complete resection should be verified by switching to blue light. Only CE marked cystoscopic equipment should be used, equipped with necessary filters to allow both standard white light cystoscopy and blue light (wavelength 380–450 nm) fluorescence cystoscopy. Children and adolescents: There is no experience of treating patients below the age of 18 years.

**CONTRAINDICATIONS** Hypersensitivity to the active substance or to any of the excipients of the solvent. Porphyria.

**WARNINGS AND PRECAUTIONS** The possibility of hypersensitivity including serious anaphylactic/ anaphylactoid reactions should always be considered. Advanced life support facilities should be readily available. Repeated use of Hexvix as part of follow-up in patients with bladder cancer has not been studied. Hexaminolevulinate should not be used in patients at high risk of bladder inflammation, e.g. after BCG therapy, or in moderate to severe leucocyturia. Widespread inflammation of the bladder should be excluded by cystoscopy before the product is administered. Inflammation may lead

to increased porphyrin build up and increased risk of local toxicity upon illumination, and false fluorescence. If a widespread inflammation in the bladder becomes evident during white light inspection, the blue light inspection should be avoided. There is an increased risk of false fluorescence in the resection area in patients who recently have undergone surgical procedures of the bladder.

**INTERACTIONS** No specific interaction studies have been performed with hexaminolevulinate.

**FERTILITY, PREGNANCY AND LACTATION** No clinical data on exposed pregnancies are available. Animal studies do not indicate direct or indirect harmful effects with respect to the reproductive toxicity. As a precautionary measure, it is preferable to avoid the use of Hexvix during pregnancy.

**UNDESIRABLE EFFECTS** Most of the reported adverse reactions from clinical studies were transient and mild or moderate in intensity. The most frequently reported adverse reactions from clinical studies were bladder spasm, reported by 2.4 % of the patients, dysuria by 1.8%, bladder pain by 1.7 % and hematuria by 1.7%, of the patients. Other commonly reported adverse reactions are: headache, nausea, vomiting, constipation, diarrhea, urinary retention, haematuria, pyrexia and post procedural pain. Uncommonly reported adverse reactions are cystitis, sepsis, urinary tract infection, insomnia, urethral pain, pollakuria, micturition urgency, urinary tract disorder, back pain, incontinence, white blood cell count increase, bilirubin and hepatic enzyme increase, post operative fever, anaemia, gout, rash and balanitis. The adverse reactions that were observed were expected, based on previous experience with standard cystoscopy and transurethral resection of the bladder (TURB) procedures.

**OVERDOSE** No case of overdose has been reported. No adverse events have been reported with prolonged instillation times exceeding 180 minutes (3 times the recommended instillation time), in one case 343 minutes. No adverse events have been reported in the dose-finding studies using twice the recommended concentration of hexaminolevulinate. There is no experience of higher light intensity than recommended or prolonged light exposure.

**INSTRUCTIONS FOR USE AND HANDLING** Hexaminolevulinate may cause sensitisation by skin contact. The product should be reconstituted under aseptic conditions using sterile equipment.

**MARKETING AUTHORISATION HOLDER**  
Photocure ASA, Hoffsvæien 4, N-0275 Oslo, Norway

**PRICE** Denmark DKK 4 718.50 Finland EUR 464.20 Norway NOK 4 234.50 Sweden SEK 4 222,00

**DATE OF REVISION OF TEXT** February 2013.

Hexvix is a registered trademark of Photocure ASA